

U. PORTO

FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2016/2017

Carlos César da Fonte Senra

Terapias Minimamente Invasivas da Artrose do Joelho

Minimally Invasive Therapies of Knee Osteoarthritis

março, 2017

FMUP

Carlos César da Fonte Senra
Terapias Minimamente Invasivas da Artrose do Joelho
Minimally Invasive Therapies of Knee Osteoarthritis

Mestrado Integrado em Medicina

Área: Medicina

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:

Doutor Carlos Vaz

Trabalho organizado de acordo com as normas da revista:

Acta Reumatológica Portuguesa

março, 2017

FMUP

Eu, Carlos César da Fonte Lima, abaixo assinado, nº mecanográfico 201108625, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 20 / 3 / 2017

Assinatura conforme cartão de identificação:

Carlos César da Fonte Lima

NOME

Carlos Gíson da Fonte Jena

NÚMERO DE ESTUDANTE

201108625

DATA DE CONCLUSÃO

20 / 03 / 2017

DESIGNAÇÃO DA ÁREA DO PROJECTO

Medicina

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

Minimally Invasive Therapies of Knee Osteoarthritis

ORIENTADOR

Doutor Carlos Vaz

COORDENADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTES TRABALHOS APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input checked="" type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTES TRABALHOS (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTES TRABALHOS.	<input type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 20/03/2017

Assinatura conforme cartão de identificação: Carlos Gíson da Fonte Jena

Aos meus pais, à minha irmã, aos meus avós,
aos meus amigos e colegas

Minimally Invasive Therapies of Knee Osteoarthritis

Abstract

Knee osteoarthritis (OA) is a very common disease and a leading cause of disability. As the knee is an easy-accessible joint, local intraarticular therapies are raising interest as new medications and medical devices are available for injections, since it can avoid the adverse effects of nonsteroidal anti-inflammatory drugs (NSAIDs) and surgery. Our review explores the clinically-evidence effective and minimally invasive treatment options available for knee OA management. The search strategy for this literature review was conducted by using the key words “knee osteoarthritis” and “intraarticular” and “therapeutics” in the PUBMED database. Among the literature, the effectiveness of Hyaluronic acid (HA) is controversial. The different molecular mass available has some responsibility in this issue. HA formulations with higher molecular mass present consistently more positive results in a long time management. Corticosteroids injections are effective in acute knee pain and effusion, without long time effect, though. Platelet-rich plasma have shown positive results among all the literature reviewed, with long time effects and disease modifying properties related to its growth factors that promote tissue healing and regeneration. A variety of novel therapies have been developed in recent years. Among them, mesenchymal stem cells injection is a promising regenerative treatment which has shown positive results in two recent randomized controlled trials, and it was considered as effective as hyaluronic acid. Genetic therapies using transduced chondrocytes expressing TGF-beta also presented promising results in two randomized trials. Polynucleotides injections also showed similar results with HA. Other single studies were found and are explored in our paper.

Introduction

Knee osteoarthritis (OA) is a very common disease that affects approximately 250 million people worldwide and is one leading cause of global disability¹. The standard recommendations of knee OA management consist of nonpharmacological approaches (such as weight loss, exercise and braces), followed by analgesic medication, including nonsteroidal anti-inflammatory drugs (NSAIDs) and lastly surgery². A recent study has suggested that surgical treatment results in greater pain relief and functional improvement compared to nonsurgical treatment, however more serious adverse events are reported in surgical treatment³. As knee is an easy-accessible joint, intraarticular therapies have an inherent appeal as it mitigates some systemic effects of NSAIDs, including gastrointestinal bleeding and myocardial infarction, and avoid adverse events of surgical treatment⁴. The purpose of our study is to evaluate minimally invasive and clinically-evidence based therapies that can be offered to those patients in which NSAIDs failed to relief pain and improve function, NSAIDs are contraindicated and are not indicated for surgery.

Methods

A search was conducted at PUBMED up to October 2016 using the key words “knee osteoarthritis” AND “intraarticular” AND “therapeutics”. Studies published prior to 2006, that involved animal experience, were related to surgical procedures/arthroscopy or that were not in English or Portuguese were excluded. An abstract quality assessment was performed by both authors to exclude low quality studies or others not related with the aim of the study.

Results

Hyaluronic Acid (“Viscossupplementation”)

Hyaluronate is a natural component of synovial fluid, acting as a joint lubricant and shock absorber. It has a molecular mass ranging from 6500 to 10900 kDa in a normal synovial fluid, however in knee OA hyaluronate is depolymerized (molecular mass 2700 to 4500 kDa) and is cleared at higher rates⁴.

17 studies were found using our inclusion/exclusion criteria evaluating clinical evidence to use hyaluronic acid (HA) injections in knee OA: 12 clinical trials (Table I) and 5 systematic reviews/meta-analysis. Among 7 randomized-controlled trials (RCT) accessed, 5 showed superiority of HA injections in pain relief and joint function⁵⁻⁹, whereas 2 showed no statistically significant difference compared to placebo^{10,11}. A meta-analysis conducted by Bannuru *et al.* evaluating 5 trials with 712 patients with knee OA suggested equal efficacy of HA injections compared to oral NSAIDs in terms of pain, function and stiffness with a better safety profile for the first option¹². In addition, three reviews concluded that HA injections could be a safe and efficacious option for knee osteoarthritis treatment¹³⁻¹⁵. On the other hand, a meta-analysis conducted by Rutjes *et al.* analysing 89 trials involving 12667 participants concluded that HA injections have a small and clinically irrelevant benefit in terms of pain management in knee OA¹⁶.

Table I – Clinical trials evaluating hyaluronic acid in knee osteoarthritis

Author/year	Study design	Conclusions
Waddell <i>et al.</i> 2015 ¹⁷	Retrospective; 50 patients requiring aspiration at the time of viscossupplementation compared with 50 matched patients without effusion	Effusion requiring aspiration at the time of hyaluronic acid injection does not negatively impact outcome
Van der Weegen <i>et al.</i> 2015 ¹⁰	RCT with 196 patients with knee OA comparing HA sodium hyaluronate with saline injections; 6 months follow-up	3 weekly injections of HA (sodium hyaluronate) were not superior to saline in 6 months follow-up
Petrella, R. J. <i>et al.</i> 2015 ¹⁸	RCT with 98 patients with knee OA comparing Hydros, Hydros-TA and cross-linked HA single injections; 26 weeks follow-up	Single injection of Hydros and Hydros-TA is effective in pain relief compared to active control; hydros-TA has a faster effect
Leighton, R. <i>et al.</i> 2014 ¹⁹	RCT with 442 patients with knee OA comparing NASHA with methylprednisolone single injection; 12 weeks follow-up	Single injection NASHA was not inferior to methylprednisolone at 12 weeks in pain score (WOMAC)

Ishijima M. et al. 2014 ²⁰	RCT with 200 patients with knee OA comparing HA injections with oral NSAID; 5 weeks follow-up	The efficacy of HA injections is not inferior to that of NSAIDs and is a more safe treatment
Arden, N. K. et al. 2014 ⁵	RCT with 218 patients with knee OA comparing NASHA with saline injections; 6 weeks follow-up	NASHA injections has a significant pain improvement at 6 weeks follow-up among patients without effusion at baseline
Strand, V. et al. 2012 ⁶	RCT with 319 patients with knee OA comparing Gel-200 with saline single injection; 13 weeks follow-up	A single injection of Gel-200 is effective and well tolerated relieving pain over 13 weeks
Navarro-Sarabia, F. et al. 2011 ⁷	RCT with 306 patients with knee OA comparing cycles of HA with saline injections; 40 months follow-up	Repeated cycles of HA injections improve pain and function in-between cycles and this beneficial carry-on for a year after the last cycle
Jorgensen, A. et al. 2010 ¹¹	RCT with 337 patients with knee OA comparing HA sodium hyaluronate with saline weekly injections for 5 weeks; 1 year follow-up	Five injections of HA (sodium hyaluronate - Hyalgan®) did not improve pain, function and analgesic consumption over 3,6,9 and 12 months follow-up
Chevalier, X. et al. 2010 ⁸	RCT with 253 patients with knee OA comparing Hylan G-F 20 with saline single injection; 26 weeks follow-up	Single injection hylan G-F 20 is safe and statistically significant in pain relief versus placebo
Altman, R. D. et al. 2009 ⁹	RCT with 588 patients with knee OA comparing HA (1% sodium hyaluronate) with placebo; 26 weeks follow-up	Three weekly injection of HA (1% sodium hyaluronate) resulted in significant pain relief and joint function over a 26 weeks follow-up
Lundsgaard, C. et al. 2008 ²¹	RCT with 251 patients to receive four weekly HA sodium hyaluronate (Hyalgan®), saline 20 ml (distension) or saline 2ml (placebo); 26 weeks follow-up	No difference between groups reducing knee pain

RCT: Randomized controlled trial; HA: hyaluronic acid; NSAID: non-steroidal anti-inflammatory drug; NASHA: cross-linked (high molecular weight) HA formulation; Gel-200: cross-linked (high molecular weight) HA formulation; WOMAC: Western Ontario and McMaster Universities Osteoarthritis Index; Hydros: cross-linked (high molecular weight) HA formulation; Hydros-TA: hydros plus 10 mg triamcinolone; Hylan G-F 20: high molecular weight (mean 6000 kDa) HA formulation;

Corticosteroids

Corticosteroids (CS) injections provides pain relief and local anti-inflammatory effect by the prostaglandin synthesis inhibition and decreasing collagenases activity. A total of 5 studies investigating the efficacy of CS injections were found. One RCT and one retrospective pilot study demonstrated that CS injections are effective in pain reduction and in functional improvement, being essential prior joint aspiration^{22,23}. CS injections are also more suitable for relieving pain in a short-time basis, with a better efficacy compared to viscosupplementation in the first 4 weeks²⁴. Another study using a different approach concluded that adding triamcinolone to hyaluronic acid improves first-

week symptoms²⁵. A RCT conducted by Bodick *et al.* reported that an extended-release formulation of triamcinolone prolongs and improve pain relief compared to standard immediate-release triamcinolone²⁶.

Platelet-rich plasma (PRP)

Eleven studies evaluating the clinical-evidence of intraarticular platelet-rich plasma (PRP) in knee OA were found: four RCT (table II), five non-randomized trials and two meta-analysis. All RCT showed that PRP injections are an effective treatment for moderate grade knee OA (level 1 of evidence in all studies)²⁷⁻³⁰, despite the different PRP formulations used. Four non-randomized, single-arm, studies, with a total of 567 patients receiving PRP injections concluded PRP to be effective in pain reduction and function improvement³¹⁻³⁴. A non-randomized trial conducted by Kon *et al.*³⁵ demonstrated that PRP is more effective reducing pain and symptoms than HA injections. A recent meta-analyse conducted by Laudy *et al.*³⁶ with ten trials comparing PRP injections with placebo revealed PRP more effective in pain reduction and function improvement at 6 months post-injection. Another meta-analyse reached the same results with PRP being an effective treatment for pain reduction compared with pre-treatment condition³⁷. Several studies revealed PRP injection being more effective in younger patients with low degree of articular degeneration^{30,31,33,35,37}.

Table II – Randomized Controlled Trials evaluating platelets-rich plasm in knee osteoarthritis

Author/year	Study design	Conclusions
Filardo, G. <i>et al.</i> 2015 ²⁷	RCT, with 192 patients with knee OA comparing PRP with HA injections; 12 months follow-up	Both treatments are equal effective improving function and reducing symptoms
Vaquerizo, V. <i>et al.</i> 2013 ²⁸	RCT, with 96 patients with knee OA comparing PRGF with HA injections; 48 weeks follow-up	PRGF proved to be better than HA injections in terms of pain reduction and functional improvement
Sanchez, M. <i>et al.</i> 2012 ²⁹	RCT, with 176 patients with knee OA comparing PRGF with HA injections; 24 weeks follow-up	PRGF showed better short-term results compared to HA injections in symptoms relief
Cerza, F. <i>et al.</i> 2012 ³⁰	RCT, with 120 patients with knee OA comparing ACP with HA injections; 24 weeks follow-up	Treatment with ACP showed a better clinical outcome compared to HA injections

RCT: randomized controlled trial; OA: osteoarthritis; HA: hyaluronic acid; PRP: platelets-rich plasma; PRGF: platelets rich in growth factors (a specific PRP formulation); ACP: autologous conditioned plasma (a specific PRP formulation).

Novel therapies

A variety of new intraarticular therapeutics has been attempted in knee OA. Our research found four recent papers related to Mesenchymal Stem Cell (MSC) therapies. A RCT with 424 patients with knee OA evaluated the relationship between cell concentration in bone marrow aspiration (without MSC isolation and expansion) and pain management, concluding that higher cell count ($>4 \times 10^8$) is associated with a better outcome³⁸. Another RCT comparing a single injection of allogenic bone marrow mesenchymal stem cells with HA in 30 randomized patients with knee OA concluded that it significantly improved cartilage quality and might be a valid alternative for pain management, lacking logistical inconveniences of autologous MSC treatment³⁹. Two phase one studies evaluating autologous adipose tissue derived MSC in knee OA patients suggested it might be a safe and promising treatment modality for these patients^{40,41}.

A RCT conducted by Lee *et al.*⁴² evaluated the efficacy of a single injection of a 1:3 mixture of genetically manipulated chondrocytes to express TGF- β 1 and normal allogenic human chondrocytes against placebo in 54 patients with knee OA concluding that in 24 weeks follow-up active treatment resulted in significantly pain and function improvement. Notwithstanding, an anaphylactic reaction to the preservation medium was reported in one patient.

Two RCT evaluating the efficacy of intraarticular injections of polynucleotides compared to HA injections in a total of 135 randomized patients with knee OA showed similar results of both treatments in pain reduction^{43,44}.

A single RCT evaluated the effect of intra-articular injections of a mixture of sodium bicarbonate with a single or double dose of calcium gluconate concluding that this treatment is effective in reducing symptoms associated with OA. A higher dose of calcium gluconate is associated with further joint-space narrowing prevention. The positive effect of bicarbonate is attributed to its alkalinity whereas calcium gluconate avoid hyperosmotic conditions in extracellular matrix by allowing the linkage between chondrocytes and bone proteins⁴⁵. Another RCT attempted 4 weekly intra-articular injection of a bisphosphonate (clodronate) resulting in a small and transient benefit over a 12 weeks follow-up compared to placebo⁴⁶. A recent prospective study found etanercept

(Tumor necrosis factor-alpha inhibitor) injections being more effective in pain management than high molecular weight HA in knee OA patients⁴⁷.

Discussion

Intra-articular HA injections in knee osteoarthritis treatment are still a controversial issue due to the contradictory results obtained from different clinical trials. Of note, two RCT that showed no difference between HA injections and placebo used a hyaluronic acid formulation with a low molecular mass (sodium hyaluronate, Hyalgan® 500 to 730 kDa)¹¹ and medium molecular mass (2200 kDa)¹⁰, respectively. Another RCT that reported no difference between HA injections, saline distension and placebo used a HA formulation with a low molecular mass (Hyalgan®)²¹. On the other hand, all studies that used hyaluronic acid formulation with a high molecular mass (mean 6000 kDa) found positive effects in knee OA^{5,6,8,18-20}. Therefore, molecular weight could represent a major issue in viscosupplementation. Previous studies had already suggested superiority of high molecular mass formulation of HA⁴⁸ and this could explain the incongruent results from several systematic reviews and meta-analysis that do not differentiate HA formulations. An effusion at baseline do not affect HA effects if we aspire it previously^{5,8,17}.

CS injections remains among the literature reviewed an efficacious option to improve joint pain and function in a short-period and fast-acting basis, so far^{22,24,25}. Some studies reinforce this short-acting improvement with no efficacy of CS injections against placebo over 12 weeks follow-up, though⁴⁹. There is not consensus regarding which CS use, doses and frequency of injection, being still a individually tailor selection⁵⁰.

PRP is an autologous blood product with a high content of growth factors stored in alpha-granules of platelets and a low content of white blood cells (WBC), which is raising interest among physicians³⁶. There are different PRP formulations with variable concentrations of growth factors and WBC³³. This variability could affect the studies outcomes since a high content of WBC, which release proinflammatory cytokines, are associated with increased pain and swelling^{51,52}. Our review were consistent with a positive effect of PRP injections in pain reduction and function improvement in knee OA. The RCTs reviewed all compared PRP with HA injections²⁷⁻³⁰. Filardo *et al.*²⁷ concluded PRP and HA injections being equal efficacious in knee OA, interestingly, the other three RCT that had positive results for PRP injections compared to HA injections used PRP formulations (Platelets rich in growth factor^{28,29} and autologous conditioned plasma³⁰) with a higher content of growth factors and a lower content of WBC. These results put in perspective the importance of PRP formulations and cell/cytokines concentration in the

final outcome. More studies are needed in this field to better understand which PRP formulation is the most effective.

MSCs therapy is a promising treatment in knee OA since those are multipotent cells with capacity to differentiate in different cellular lines of mesodermal origin including cartilage, ligaments and tendon³⁸. MSC could be autologous: bone marrow derived or adipose tissue derived with inconvenience related to the invasive bone marrow aspirations and adipose tissue extraction and also, MSC isolation and cell culture not suitable for an outpatient setting. In alternative other authors instead of isolating MSC, use bone marrow concentrates containing a fraction of MSC and other bone marrow cells (including hematopoietic stem cells, monocyte precursor cells, macrophages, T cells, B cells and others) with comparable results without MSC being isolated, suggesting that other bone marrow cells could affect tissue healing³⁸. Adipose-derived MSC are a promising alternative of autologous MSC therapies but further RCT studies are needed to access efficacy⁴¹. Allogeneic MSC are a cheaper and more convenient way for MSC intraarticular therapies. Despite the possibility of host immune rejection, MSC are *immune evasive* and inhibit immune responses. A RCT with 15 patients receiving allogeneic MSC reported no safety concerns³⁹. In spite of the potential of allogeneic MSC, larger RCT are needed to evaluate safety profile.

Genetically manipulated chondrocytes expressing TGF- β are a recent treatment modality for knee OA. TGF- β is thought to have regenerative, anti-inflammatory and immunosuppressive properties, stimulating proteoglycan synthesis and growth of articular chondrocytes. This cell-mediated gene therapy consist of a mixture of transduced allogeneic chondrocytes overexpressing TGF- β and normal allogeneic chondrocytes. Non-transduced chondrocytes are included to fill cartilage defects and act as an additional target for TGF- β signalling⁵³. This therapy had promising results in a RCT and a phase two study, being sufficient for further clinical testing^{42,53}.

Intraarticular polynucleotides act as a 3D gel with a high content of water with mechanical effects in joint lubrication and moisturizing. In addition, polynucleotides provide synovial fluid with nucleotides, nucleosides, purine and pyrimidine that can therefore support the physiological repair processes of cartilage. Two RCT showed similar results of intraarticular polynucleotides compared to HA injections, being an alternative for this last treatment ^{43,44}.

Intra-articular etanercept had promising results in knee OA pain management, suggesting TNF- α is one factor that induces OA pain. However, one single trial was found and more studies are needed to evaluate efficacy and safety⁴⁷.

References

1. Cross M, Smith E, Hoy D, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014;73:1323-30.
2. Hochberg MC, Altman RD, April KT, et al. American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)* 2012;64:465-74.
3. Skou ST, Roos EM, Laursen MB, et al. A Randomized, Controlled Trial of Total Knee Replacement. *N Engl J Med* 2015;373:1597-606.
4. Hunter DJ. Viscosupplementation for osteoarthritis of the knee. *N Engl J Med* 2015;372:1040-7.
5. Arden NK, Akermark C, Andersson M, Todman MG, Altman RD. A randomized saline-controlled trial of NASHA hyaluronic acid for knee osteoarthritis. *Curr Med Res Opin* 2014;30:279-86.
6. Strand V, Baraf HS, Lavin PT, Lim S, Hosokawa H. A multicenter, randomized controlled trial comparing a single intra-articular injection of Gel-200, a new cross-linked formulation of hyaluronic acid, to phosphate buffered saline for treatment of osteoarthritis of the knee. *Osteoarthritis Cartilage* 2012;20:350-6.
7. Navarro-Sarabia F, Coronel P, Collantes E, et al. A 40-month multicentre, randomised placebo-controlled study to assess the efficacy and carry-over effect of repeated intra-articular injections of hyaluronic acid in knee osteoarthritis: the AMELIA project. *Ann Rheum Dis* 2011;70:1957-62.
8. Chevalier X, Jerosch J, Goupille P, et al. Single, intra-articular treatment with 6 ml hylan G-F 20 in patients with symptomatic primary osteoarthritis of the knee: a randomised, multicentre, double-blind, placebo controlled trial. *Ann Rheum Dis* 2010;69:113-9.
9. Altman RD, Rosen JE, Bloch DA, Hatoum HT, Korner P. A double-blind, randomized, saline-controlled study of the efficacy and safety of EUFLEXXA for treatment of painful osteoarthritis of the knee, with an open-label safety extension (the FLEXX trial). *Semin Arthritis Rheum* 2009;39:1-9.
10. van der Weegen W, Wullems JA, Bos E, Noten H, van Drumpt RA. No difference between intra-articular injection of hyaluronic acid and placebo for mild to moderate knee osteoarthritis: a randomized, controlled, double-blind trial. *J Arthroplasty* 2015;30:754-7.
11. Jorgensen A, Stengaard-Pedersen K, Simonsen O, et al. Intra-articular hyaluronan is without clinical effect in knee osteoarthritis: a multicentre, randomised, placebo-controlled, double-blind study of 337 patients followed for 1 year. *Ann Rheum Dis* 2010;69:1097-102.
12. Bannuru RR, Vaysbrot EE, Sullivan MC, McAlindon TE. Relative efficacy of hyaluronic acid in comparison with NSAIDs for knee osteoarthritis: a systematic review and meta-analysis. *Semin Arthritis Rheum* 2014;43:593-9.
13. Trigkilidas D, Anand A. The effectiveness of hyaluronic acid intra-articular injections in managing osteoarthritic knee pain. *Ann R Coll Surg Engl* 2013;95:545-51.
14. Reid MC. Viscosupplementation for osteoarthritis: a primer for primary care physicians. *Adv Ther* 2013;30:967-86.
15. Ray TR. Using viscosupplementation to treat knee osteoarthritis. *Phys Sportsmed* 2013;41:16-24.

16. Rutjes AW, Juni P, da Costa BR, Trelle S, Nuesch E, Reichenbach S. Viscosupplementation for osteoarthritis of the knee: a systematic review and meta-analysis. *Ann Intern Med* 2012;157:180-91.
17. Waddell BS, Waddell WH, Waddell DD. Comparison of Efficacy and Tolerability of Hylan G-F 20 in Patients with and without Effusions at the Time of Initial Injection. *J Knee Surg* 2015;28:213-22.
18. Petrella RJ, Emans PJ, Alleyne J, Dellaert F, Gill DP, Maroney M. Safety and performance of Hydros and Hydros-TA for knee osteoarthritis: a prospective, multicenter, randomized, double-blind feasibility trial. *BMC Musculoskelet Disord* 2015;16:57.
19. Leighton R, Akermark C, Therrien R, et al. NASHA hyaluronic acid vs. methylprednisolone for knee osteoarthritis: a prospective, multi-centre, randomized, non-inferiority trial. *Osteoarthritis Cartilage* 2014;22:17-25.
20. Ishijima M, Nakamura T, Shimizu K, et al. Intra-articular hyaluronic acid injection versus oral non-steroidal anti-inflammatory drug for the treatment of knee osteoarthritis: a multi-center, randomized, open-label, non-inferiority trial. *Arthritis Res Ther* 2014;16:R18.
21. Lundsgaard C, Dufour N, Fallentin E, Winkel P, Gluud C. Intra-articular sodium hyaluronate 2 mL versus physiological saline 20 mL versus physiological saline 2 mL for painful knee osteoarthritis: a randomized clinical trial. *Scand J Rheumatol* 2008;37:142-50.
22. Yavuz U, Sokucu S, Albayrak A, Ozturk K. Efficacy comparisons of the intraarticular steroidal agents in the patients with knee osteoarthritis. *Rheumatol Int* 2012;32:3391-6.
23. Leung A, Liew D, Lim J, et al. The effect of joint aspiration and corticosteroid injections in osteoarthritis of the knee. *Int J Rheum Dis* 2011;14:384-9.
24. Bannuru RR, Natov NS, Obadan IE, Price LL, Schmid CH, McAlindon TE. Therapeutic trajectory of hyaluronic acid versus corticosteroids in the treatment of knee osteoarthritis: a systematic review and meta-analysis. *Arthritis Rheum* 2009;61:1704-11.
25. de Campos GC, Rezende MU, Pailo AF, Frucchi R, Camargo OP. Adding triamcinolone improves viscosupplementation: a randomized clinical trial. *Clin Orthop Relat Res* 2013;471:613-20.
26. Bodick N, Lufkin J, Willwerth C, et al. An intra-articular, extended-release formulation of triamcinolone acetonide prolongs and amplifies analgesic effect in patients with osteoarthritis of the knee: a randomized clinical trial. *J Bone Joint Surg Am* 2015;97:877-88.
27. Filardo G, Di Matteo B, Di Martino A, et al. Platelet-Rich Plasma Intra-articular Knee Injections Show No Superiority Versus Viscosupplementation: A Randomized Controlled Trial. *Am J Sports Med* 2015;43:1575-82.
28. Vaquerizo V, Plasencia MA, Arribas I, et al. Comparison of intra-articular injections of plasma rich in growth factors (PRGF-Endoret) versus Durolane hyaluronic acid in the treatment of patients with symptomatic osteoarthritis: a randomized controlled trial. *Arthroscopy* 2013;29:1635-43.
29. Sanchez M, Fiz N, Azofra J, et al. A randomized clinical trial evaluating plasma rich in growth factors (PRGF-Endoret) versus hyaluronic acid in the short-term treatment of symptomatic knee osteoarthritis. *Arthroscopy* 2012;28:1070-8.
30. Cerza F, Carni S, Carcangiu A, et al. Comparison between hyaluronic acid and platelet-rich plasma, intra-articular infiltration in the treatment of gonarthrosis. *Am J Sports Med* 2012;40:2822-7.

31. Jang SJ, Kim JD, Cha SS. Platelet-rich plasma (PRP) injections as an effective treatment for early osteoarthritis. *Eur J Orthop Surg Traumatol* 2013;23:573-80.
32. Wang-Saegusa A, Cugat R, Ares O, Seijas R, Cusco X, Garcia-Balletbo M. Infiltration of plasma rich in growth factors for osteoarthritis of the knee short-term effects on function and quality of life. *Arch Orthop Trauma Surg* 2011;131:311-7.
33. Kon E, Buda R, Filardo G, et al. Platelet-rich plasma: intra-articular knee injections produced favorable results on degenerative cartilage lesions. *Knee Surg Sports Traumatol Arthrosc* 2010;18:472-9.
34. Filardo G, Kon E, Buda R, et al. Platelet-rich plasma intra-articular knee injections for the treatment of degenerative cartilage lesions and osteoarthritis. *Knee Surg Sports Traumatol Arthrosc* 2011;19:528-35.
35. Kon E, Mandelbaum B, Buda R, et al. Platelet-rich plasma intra-articular injection versus hyaluronic acid viscosupplementation as treatments for cartilage pathology: from early degeneration to osteoarthritis. *Arthroscopy* 2011;27:1490-501.
36. Laudy AB, Bakker EW, Rekers M, Moen MH. Efficacy of platelet-rich plasma injections in osteoarthritis of the knee: a systematic review and meta-analysis. *Br J Sports Med* 2015;49:657-72.
37. Chang KV, Hung CY, Aliwarga F, Wang TG, Han DS, Chen WS. Comparative effectiveness of platelet-rich plasma injections for treating knee joint cartilage degenerative pathology: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2014;95:562-75.
38. Centeno CJ, Al-Sayegh H, Bashir J, Goodyear S, Freeman MD. A dose response analysis of a specific bone marrow concentrate treatment protocol for knee osteoarthritis. *BMC Musculoskelet Disord* 2015;16:258.
39. Vega A, Martin-Ferrero MA, Del Canto F, et al. Treatment of Knee Osteoarthritis With Allogeneic Bone Marrow Mesenchymal Stem Cells: A Randomized Controlled Trial. *Transplantation* 2015;99:1681-90.
40. Jo CH, Lee YG, Shin WH, et al. Intra-articular injection of mesenchymal stem cells for the treatment of osteoarthritis of the knee: a proof-of-concept clinical trial. *Stem Cells* 2014;32:1254-66.
41. Pers YM, Rackwitz L, Ferreira R, et al. Adipose Mesenchymal Stromal Cell-Based Therapy for Severe Osteoarthritis of the Knee: A Phase I Dose-Escalation Trial. *Stem cells translational medicine* 2016;5:847-56.
42. Lee MC, Ha CW, Elmallah RK, et al. A placebo-controlled randomised trial to assess the effect of TGF- α 1-expressing chondrocytes in patients with arthritis of the knee. *Bone Joint J* 2015;97-B:924-32.
43. Giarratana LS, Marelli BM, Crapanzano C, et al. A randomized double-blind clinical trial on the treatment of knee osteoarthritis: the efficacy of polynucleotides compared to standard hyaluronan viscosupplementation. *Knee* 2014;21:661-8.
44. Vanelli R, Costa P, Rossi SM, Benazzo F. Efficacy of intra-articular polynucleotides in the treatment of knee osteoarthritis: a randomized, double-blind clinical trial. *Knee Surg Sports Traumatol Arthrosc* 2010;18:901-7.
45. Garcia-Padilla S, Duarte-Vazquez MA, Gonzalez-Romero KE, Caamano Mdel C, Rosado JL. Effectiveness of intra-articular injections of sodium bicarbonate and calcium gluconate in the treatment of osteoarthritis of the knee: a randomized double-blind clinical trial. *BMC Musculoskelet Disord* 2015;16:114.

46. Rossini M, Adami S, Fracassi E, et al. Effects of intra-articular clodronate in the treatment of knee osteoarthritis: results of a double-blind, randomized placebo-controlled trial. *Rheumatol Int* 2015;35:255-63.
47. Ohtori S, Orita S, Yamauchi K, et al. Efficacy of Direct Injection of Etanercept into Knee Joints for Pain in Moderate and Severe Knee Osteoarthritis. *Yonsei Med J* 2015;56:1379-83.
48. Raman R, Dutta A, Day N, Sharma HK, Shaw CJ, Johnson GV. Efficacy of Hylan G-F 20 and Sodium Hyaluronate in the treatment of osteoarthritis of the knee -- a prospective randomized clinical trial. *Knee* 2008;15:318-24.
49. Henriksen M, Christensen R, Klokke L, et al. Evaluation of the benefit of corticosteroid injection before exercise therapy in patients with osteoarthritis of the knee: a randomized clinical trial. *JAMA Intern Med* 2015;175:923-30.
50. Douglas RJ. Corticosteroid injection into the osteoarthritic knee: drug selection, dose, and injection frequency. *Int J Clin Pract* 2012;66:699-704.
51. Pourcho AM, Smith J, Wisniewski SJ, Sellon JL. Intraarticular platelet-rich plasma injection in the treatment of knee osteoarthritis: review and recommendations. *Am J Phys Med Rehabil* 2014;93:S108-21.
52. Filardo G, Kon E, Pereira Ruiz MT, et al. Platelet-rich plasma intra-articular injections for cartilage degeneration and osteoarthritis: single- versus double-spinning approach. *Knee Surg Sports Traumatol Arthrosc* 2012;20:2082-91.
53. Ha CW, Cho JJ, Elmallah RK, et al. A Multicenter, Single-Blind, Phase IIa Clinical Trial to Evaluate the Efficacy and Safety of a Cell-Mediated Gene Therapy in Degenerative Knee Arthritis Patients. *Hum Gene Ther Clin Dev* 2015;26:125-30.

Agradecimentos

Ao Professor Doutor Carlos Vaz pela sua orientação, total apoio, disponibilidade, pelo saber que transmitiu, pelas opiniões e críticas, total colaboração no solucionar de problemas e dúvidas e por todas as palavras de incentivo.

À minha irmã, pela revisão linguística.

Aos meus colegas e amigos por todo o apoio e incentivo.

Acta Reumatológica Portuguesa

Normas de publicação

Objectivos e âmbito

A Acta Reumatológica Portuguesa (ARP) é uma publicação científica internacional, revista por pares, abrangendo aspectos clínicos e experimentais das doenças reumáticas. São publicados artigos originais, artigos de revisão, casos clínicos, imagens em Reumatologia, cartas ao editor e artigos que visam melhorar a Prática Clínica (recomendações e protocolos clínicos, por exemplo).

A ARP foi fundada em 1973 como órgão científico oficial da Sociedade Portuguesa de Reumatologia e subscrive os requisitos para apresentação de artigos a revistas biomédicas elaboradas pela Comissão Internacional de Editores de Revistas Médicas (International Committee of Medical Journal Editors), publicada na íntegra inicialmente em N Engl J Med 1991; 324: 42428 e actualizada em Outubro de 2008 e disponível em www.ICMJE.org. A política editorial da ARP segue as Recomendações de Política Editorial (Editorial Policy Statements) emitidas pelo Conselho de Editores Científicos (Council of Science Editors), disponíveis em http://www.councilscienceeditors.org/files/public/entire_whitepaper.pdf.

A ARP publica preferencialmente artigos escritos na língua Inglesa. Caso os autores optem por submeter em Português, poderá ser solicitada a sua tradução para a língua Inglesa. O rigor e a exactidão dos conteúdos, assim como as opiniões expressas, são da exclusiva responsabilidade do(s) autor(es).

Os artigos submetidos devem ser originais e não podem ter sido publicados previamente.

Os artigos publicados constituirão propriedade da revista, não podendo ser reproduzidos, no seu todo ou em parte, sem a prévia autorização dos editores.

A Revista está indexada no PubMed/Medline e os seus artigos estão disponíveis online na íntegra, com acesso aberto e gratuito.

Instruções para submissão de artigos

A Acta Reumatológica Portuguesa oferece aos autores um sistema de submissão e revisão de artigos a funcionar exclusivamente online. Acedendo ao website da ARP (www.actareumatologica.pt), os autores poderão submeter os seus artigos e acompanhar o seu estado no processo de revisão. Os autores serão notificados por email no próprio dia em que o(s) seu(s) artigo(s) sofra(m) alterações relevantes durante o processo editorial.

De modo a submeter um manuscrito, os autores deverão **criar uma conta de utilizador**:

- Aceder ao website da ARP (www.actareumatologica.pt), clicar no link "Entrar" seguido de "Registo" e seguir cuidadosamente todas as instruções fornecidas. Um email de activação será enviado para a sua conta de email. Para activar a conta ARP é necessário seguir o link fornecido no corpo desse email, que automaticamente o redirecionará para uma mensagem de registo no website da ARP.

Após a criação de uma conta ARP, os autores poderão **submeter e acompanhar o progresso do(s) seu(s) artigo(s)**:

- Aceder ao website da ARP (www.actareumatologica.pt), entrar na area privada e clicar no link "Submeter artigo". Preencher o formulário seguindo as instruções cuidadosamente.

Instruções aos autores

Para evitar atrasos no processo de revisão, leia cuidadosamente as instruções e assegure-se de que o seu manuscrito está de acordo com os requisitos da ARP antes de submeter (número de palavras e formato).

- **Título do artigo:** o título deve descrever brevemente o conteúdo do artigo. Não devem ser usadas abreviaturas. Deve ser indicado um curto título para rodapé. Nos artigos escritos em português é necessário incluir o título em inglês.
- **Nome dos autores e afiliações**
- **Informações do autor responsável pela correspondência:** nome, morada, telefone e endereço electrónico
- **Resumo:** com um máximo de 350 palavras ,deve incluir objectivos, material e métodos, resultados e conclusões. Para os casos clínicos o limite de palavras é 180.

Tipos de artigo:

- **Editoriais:** Os Editoriais serão solicitados por convite do Editor e constituirão comentários sobre tópicos actuais ou sobre artigos publicados na revista. O texto dos Editoriais não deverá exceder as 1200 palavras, um máximo de 15 referências e pode conter uma figura ou tabela.
- **Artigos de Revisão:** Preferencialmente, os Artigos de Revisão serão também solicitados pelo Editor. No entanto, os autores interessados em apresentar um Artigo de Revisão podem contactar o Editor para discussão dos temas a apresentar no artigo, o qual não deverá exceder as 4000 palavras, 6 Tabelas ou Figuras e 100 referências;
- **Artigos Originais:** O texto dos Artigos Originais deve ser apresentado com uma Introdução, Material e Métodos, Resultados, Discussão e Conclusão. Não deverá exceder as 4000 palavras, 6 Tabelas ou Figuras e 60 referências;
- **Prática Clínica:** O texto dos artigos de Prática Clínica deve ser apresentado com uma Introdução, Material e Métodos, Resultados, Discussão e Conclusão. Não deverá exceder as 4000 palavras, 6 Tabelas ou Figuras e 60 referências;
- **Casos Clínicos:** O texto dos Casos Clínicos deverá ser apresentado com uma Introdução, Caso Clínico e Discussão. Os Casos Clínicos deverão ser acompanhados de figuras ilustrativas/tabelas (máximo de 6), não deverão exceder as 2000 palavras e 25 referências;
- **Imagens em Reumatologia:** Imagens representando manifestações clínicas raras ou de particular interesse podem ser submetidas (no máximo 4). O texto acompanhante não deverá exceder as 500 palavras e 5 referências
- **Cartas ao Editor:** As Cartas ao Editor deverão constituir um comentário critico a um artigo da Revista ou uma pequena nota sobre um tema ou caso clínico. O texto não deverá exceder as 600 palavras, uma Figura/Tabela e um máximo de 10 referências.

Tabelas: As Tabelas a inserir devem ser assinaladas no texto em numeração romana. Cada Tabela deverá possuir um título e não deverá apresentar linhas verticais. As linhas horizontais só deverão ser usadas como separadores de título e subtítulos. Todas as abreviaturas usadas devem ser explicadas na parte inferior da Tabela.

Figuras: As Figuras a inserir devem ser assinaladas no texto em numeração árabe e apresentar legendas. Cada Figura deve ser importada individualmente em format JPEG ou TIFF de alta qualidade. O Editor reserva o direito de agrupar Figuras ou alterar o seu tamanho de modo a rentabilizar o uso da páginas.

Referências: As Referências bibliográficas devem ser classificadas e numeradas por ordem de entrada no texto e em superscript. As abreviaturas usadas na nomeação das revistas devem ser as utilizadas pelo Index Medicus. Nas Referências com 6 ou menos autores, todos devem ser nomeados. Nas Referências com 7 ou mais autores, devem ser nomeados os 3 primeiros seguidos de et al. Os números de página inicial e final devem ser totalmente apresentados (ex. 565-569 e não 565-9). Não indicar o número da Revista nem o mês da publicação. As Referências correspondentes a trabalhos não publicados, apresentações ou observações pessoais, devem ser inseridas no próprio texto (em parenthesis) e não como Referências convencionais. Os autores são responsáveis pela exactidão das Referências apresentadas. Seguem-se alguns exemplos de como devem constar os vários tipos de Referências:

Revista

Nome(s) e iniciais do(s) autor(es). Título do artigo. Nome da Revista Ano; Volume: Página(s).

Ex: Hill J, Bird HA, Hopkins R, Lawton C, Wright C. Survey of satisfaction with care in a rheumatology outpatient clinic. Ann Rheum Dis 1992; 51: 195-197.

Artigo publicado online (inserir DOI)

Nome(s) e iniciais do(s) autor(es). Título do artigo. Nome da Revista Published Online First: data. doi.

Ex: Merkel PA, Curthbertson D, Hellmich B et al. Comparison of disease activity measures for ANCA-associated vasculitis. Ann Rheum Dis Published Online First: 29 July 2008. doi:10.1136/ard.2008.097758.

Capítulo de livro

Nome(s) e iniciais do(s) autor(es) do capítulo. Título do capítulo. In: Nome(s) e iniciais do(s) editor(es) medico(s). Título do livro. Cidade: Nome da casa editorial, ano de publicação: primeira a última página do capítulo.

Ex: Stewart AF. Hypercalcemia resulting from medications. In: Favus MD, ed Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism. New York: Raven Press, 1991: 177-178.

Livro

Nome(s) e iniciais do(s) autor(es). Título do livro. Cidade: Nome da casa editorial, ano de publicação: página(s).

Ex: Lorig K. Patient Education. A practical approach. St Louis: Mosby-Year Book, 1992: 51.

Documento electrónico

Título do documento. <http://address>. Data de acesso.

Ex: Programa Nacional de Luta Contra a Tuberculose. Sistema de Vigilância (SVIG-TB).

Direcção-Geral da Saúde – Divisão de Doenças Transmissíveis, Março de 2005.

<http://www.dgsaude.pt/upload/membro.id/ficheiros/io06875.pdf>. Accessed in January 25th 2008.

Agradecimentos

Incluir nesta secção agradecimentos a pessoas que tenham contribuído para o trabalho mas sem autoria. Instituições ou fontes de apoio financeiro também poderão aqui ser indicadas.

Processo de Revisão

Os artigos submetidos são enviados a revisores especializados no tema do artigo. Concluída a revisão do artigo, os autores são notificados, recebendo os pareceres e comentários dos revisores acerca do estado do mesmo. Com base nesses pareceres, os autores deverão editar o artigo, corrigi-lo e resubmetê-lo para nova revisão. Na resubmissão de um artigo terão de ser incluídas em anexo uma carta-resposta aos Revisores e um .doc com uma versão do artigo em *track changes*. Este documento não deverá ter qualquer identificação dos autores nem as respectivas afiliações. Caso o artigo se mantenha sem uma resposta dos autores durante mais de 6 meses, a Equipa editorial reserva-se no direito de o retirar do processo de revisão.

CrITÉrios de Revisão

Os critérios de aceitação de um artigo para publicação têm em consideração a qualidade e originalidade do artigo apresentado, a excelência na redacção e organização do mesmo e o potencial impacto na literatura médica.

Todos os artigos aceites para publicação serão mantidos e apresentados como “*online-first*” até que os Editores os seleccionem para integrar um número específico da Revista.

Revisão de provas

Os autores dos artigos aceites para publicação receberão uma versão digital da prova do artigo para valiação. Assim, as provas devem ser revistas durante as 72h que sucedem a sua recepção. Os autores devem verificar o texto, Figuras, Tabelas, Legendas e Referências, e contactar os Editores no caso em que sejam necessárias alterações. Apenas pequenas alterações e correcções tipográficas são permitidas nesta fase.

Direitos de autor

Após aceitação para publicação, os autores transferem para a ARP os direitos de autor do manuscrito.